

Physics 101H

General Physics 1 - Honors




Lecture 9 - 9/15/22

2D motion




What was the most important equation we discussed yesterday?

Example: A wheel of radius R is stuck in the mud, spinning in place with the rim moving at speed v . Bits of the mud depart from the wheel at various random locations, as in the figure. Find the maximum height that the mud reaches, assuming $v^2 > gR$.



Example: An elephant is located on Earth's surface at a latitude λ . Calculate the centripetal acceleration of the elephant resulting from the rotation of the Earth about its polar axis.





Is motion in a vertical circle uniform circular motion or nonuniform circular motion?



Summary

Topics

Yesterday: kinematics in 2D [chapter 4]

- Examples in 2D
- Uniform circular motion

Today: kinematics in 2D [chapter 4]

- Circular motion
- Relative velocity
- Galilean transformations

Announcements

Yesterday: Problem set 2 assigned

Nonuniform motion



Generalise our decomposition of the acceleration to a more general case:

- motion along a curved path with variable speed

Acceleration can be broken into **radial** and **tangential acceleration**



Relative motion

Motion gets even more interesting

Relative velocity



Position is defined by first choosing a specific reference frame

Velocity is therefore defined with respect to a specific reference frame

So how do we compare observations made by experimenters in different reference frames that are moving with respect to each other?

Frame transformations



Galilean transformations – relate observations in moving reference frames
Transformations relevant to **nonrelativistic motion** (slowly moving objects)

Lorentz transformations – also relate observations in moving reference frames!
Transformations relevant to **special relativity**

- ⦿ Describes fast moving objects
- ⦿ Entangle both space and time (i.e. **spacetime**), not just space
- ⦿ Ensure objects with mass cannot travel faster than light in a vacuum

Want more practice?



Try the following problems **Chapter 4** of the [textbook](#):

- Conceptual questions: 5, 7, 11, 15
- 2D kinematics as vectors: 19, 23, 27, 31
- Projectile motion: 33, 35, 39, 43, 51, **99**
- Uniform circular motion: 61, 63, 67, **87**
- Relative motion: 73, 77

Answers are provided for questions with **blue** numbers (odd numbered)

Click on the number to be taken to the answer.

But make sure you at least **try** the problem first!

49. A fluid ounce is about 30 mL. What is the volume of a 12 fl-oz can of soda pop in cubic meters?

1.4 Dimensional Analysis

50. A student is trying to remember some formulas from geometry. In what follows, assume A is area, V is volume, and all other variables are lengths. Determine which formulas are dimensionally consistent. (a) $V = \pi r^2 h$; (b) $A = 2\pi r^2 + 2\pi r h$; (c) $V = 0.5bh$; (d) $V = \pi d^2$; (e) $V = \pi d^3/6$.

51. Consider the physical quantities s , v , a , and t with dimensions $[s] = L$, $[v] = LT^{-1}$, $[a] = LT^{-2}$, and $[t] = T$. Determine whether each of the following equations is dimensionally consistent. (a) $v^2 = 2as$; (b) $s = vt^2 + 0.5at^2$; (c) $v = st$; (d) $a = vt$.

52. Consider the physical quantities m , s , v , a , and t with dimensions $[m] = M$, $[s] = L$, $[v] = LT^{-1}$, $[a] = LT^{-2}$, and $[t] = T$. Assuming each of the following equations is dimensionally consistent, find the dimension of the quantity on the left-hand side of the equation: (a) $F = ma$; (b) $K = 0.5mv^2$; (c) $p = mv$; (d) $W = mas$; (e) $L = mvr$.

53. Suppose quantity s is a length and quantity t is a time. Suppose the quantities v and a are defined by $v = ds/dt$ and $a = dv/dt$. (a) What is the dimension of v ? (b) What is the dimension of the quantity a ? What are the dimensions of (c) $\int v dt$, (d) $\int a dt$, and (e) da/dt ?

54. Suppose $[V] = L^3$, $[\rho] = ML^{-3}$, and $[t] = T$. (a) What is the dimension of $\int \rho dV$? (b) What is the dimension of dV/dt ? (c) What is the dimension of $\rho(dV/dt)$?

55. The arc length formula says the length s of arc subtended by angle Θ in a circle of radius r is given by the equation $s = r\Theta$. What are the dimensions of (a) s , (b) r , and (c) Θ ?

1.5 Estimates and Fermi Calculations

56. Assuming the human body is made primarily of water, estimate the volume of a person.



Summary

Topics

Today: kinematics in 2D [chapter 4]

- Circular motion
- Relative velocity
- Galilean transformations

Tomorrow: Forces [chapter 5]

- Types of forces
- Field forces and contact forces

Announcements

Yesterday: Problem set 2 assigned

**NEXT WEEK:
NO CLASS ON THURSDAY OR FRIDAY**




PHYSICS 101 - HONORS

Lecture 9 9/15/22

Wheel example: (slides 3 and 4)

First note:



$$v_x = v \sin \theta$$
$$v_y = v \cos \theta$$

Recall $h = \frac{v_i^2 \sin^2 \theta}{2g}$ but for θ defined by

$$\text{So here } h = \frac{v^2 \sin^2(90 - \theta)}{2g} = \frac{v^2 \cos^2 \theta}{2g}$$

\Rightarrow total height is initial height + height gained

$$H = (R + R \sin \theta) + \frac{v^2 \cos^2 \theta}{2g}$$

To maximise, take $\frac{dH}{d\theta} = 0$

$$\Rightarrow \frac{dH}{d\theta} = R \cos \theta - \frac{v^2}{g} \sin \theta \cos \theta = 0$$

$$\Rightarrow \left(R - \frac{v^2}{g} \sin \theta \right) \cos \theta = 0$$

$$\sin \theta = \frac{gR}{v^2} \quad \theta = \pi/2 \quad (-\pi/2 \text{ unphysical})$$

this is < 1
if $v^2 > gR$ ✓


$$\text{So } H_{\max} = R + R \cdot \left(\frac{gR}{v^2} \right) + \frac{v^2}{2g} \left(1 - \frac{g^2 R^2}{v^4} \right) = R + \frac{gR^2}{v^2} + \frac{v^2}{2g} - \frac{gR^2}{2v^2}$$

$$= \left[R + \frac{gR^2}{2v^2} + \frac{v^2}{2g} \right]$$

Rotating elephant example

Distance from the polar axis at latitude λ is

$$r = R_E \cos \lambda$$




The period of rotation is T_E (ie 24 hrs)
so the ω of the elephant is

$$v = \frac{2\pi r}{T_E} = \frac{2\pi R_E \cos \lambda}{T_E}$$


And the acceleration is therefore

$$a_c = \frac{v^2}{r} = \left(\frac{4\pi^2 R_E^2 \cos^2 \lambda}{T_E^2} \right) / R_E \cos \lambda = \boxed{\frac{4\pi^2 R_E \cos \lambda}{T_E^2}}$$

Nonuniform motion (slide 6)



Nonuniform circular motion



- velocity still tangent to path
- but \bar{a} not necessarily \perp to \bar{v}
- still decompose \bar{a} as

$$\bar{a} = \bar{a}_r + \bar{a}_T$$

radial,
 \perp to \bar{v}
↑
tangential
 \parallel to \bar{v}


$$|\bar{a}| = \sqrt{|\bar{a}_r|^2 + |\bar{a}_T|^2}$$

Note $\bar{a} = \bar{a}(t)$
so decomposition changes!

$$|\bar{a}_T| = \left| \frac{d\bar{v}}{dt} \right| \rightarrow \text{changing speed}$$

$$|\bar{a}_r| = a_c = \frac{v^2}{r} \rightarrow \text{changing direction}$$

Relative velocity (slide 8)



$$\bar{r}_b = \bar{r}_a + \bar{r}_{AB} \quad \text{but if object is moving?}$$

$$\Rightarrow \bar{u}_b = \frac{d\bar{r}_b}{dt}$$

if A and B not moving ($\frac{d\bar{r}_{AB}}{dt} = 0$) then

$$\bar{u}_b = \frac{d\bar{r}_b}{dt} = \frac{d\bar{r}_a}{dt}$$

what happens if the frames are moving?

Galilean transformations (slide 9)

Suppose A and B coincide at time $t = 0 \Rightarrow \vec{r}_{AB} = \vec{v}_{AB}t$

$$\text{Now } \frac{d}{dt} \vec{r}_b = \frac{d}{dt} (\vec{r}_a + \vec{r}_{AB}) = \frac{d\vec{r}_a}{dt} + \frac{d}{dt} (\vec{v}_{AB}t) = \vec{u}_a + \vec{v}_{AB}$$

$$\Rightarrow \boxed{\vec{u}_b = \vec{u}_a + \vec{v}_{AB}}$$

assuming \vec{v}_{AB} constant,
only holds for inertial
(nonaccelerating frames)!

Velocities in frames moving
at constant velocity with
respect to each other add!

$$\text{Note that } \vec{a}_b = \frac{d\vec{u}_b}{dt} = \frac{d\vec{u}_a}{dt} + \frac{d\vec{v}_{AB}}{dt} = \frac{d\vec{u}_a}{dt}$$

Accelerations in frames moving at constant velocity
with respect to each other are the same!